

# Horticulture 2016 Newsletter

No. 8 February 23, 2016

2021 Throckmorton Plant Science Cntr.  
Manhattan, KS 66506 (785) 532-6173

---

**Video of the Week:** [Soil Testing: Using Test Strips](#)

## FRUIT

### Growing Blueberries



Blueberries are not native to Kansas but will grow in the eastern half of the state with good preparation. They are related to azaleas and rhododendrons and require an acid pH, preferably 4.8 to 5.2. Blueberries do not have root hairs, so watering and mulching are important.

It is best to start planting preparations a year ahead of time to allow for pH adjustment, weed control, and the addition of organic matter. The first step is a soil test to determine how much the pH needs to

be reduced. For a pH up to 5.5, the addition of sphagnum peat moss at the rate of 2 cubic feet per 100 square feet will be adequate. For a pH 5.5 to 6.0, add 1 pound of sulfur per 100 square feet of bed in addition to the peat moss. For a pH 6.0 to 6.5, add 1.5 pounds of sulfur per 100 square feet of bed. For pH levels above 6.5, use 2 pounds of sulfur per 100 square feet of bed and double the amount of sphagnum peat moss suggested earlier. Do not use aluminum sulfate to correct a high pH because excessive levels of aluminum can be toxic to blueberries. For each 0.5 movement up the pH scale from 6.5, add an additional pound of sulfur. Sulfur can be applied as a dust, but pelletized sulfur is much easier to spread.

Treat only the row. Row width should be 8 feet. Blueberries are normally spaced about 5 feet within the row. Sulfur takes time to react, so allow as much time as possible between sulfur application and planting. Blueberries will bear more if you plant more than one variety.

Recommended varieties vary, but you may want to try Bluecrop because it is adaptable. Patriot also seems to do well. You may want to try some other varieties.

Blueberries should be mulched. Sawdust is the traditional material, but straw and wood chips will work as well. Mulch to a depth of about 3 inches.

Blueberries must be irrigated. Soils should be kept moist but never waterlogged. Adding peat moss to the planting row will elevate the planting bed enough that standing water should not be

an issue. An elevated bed will dry out more quickly, so there must be a means of adding water. Trickle irrigation works well. Watering twice a week during the summer with enough water to wet the soil 8 inches deep should be sufficient except under extreme heat. Watering once a week may be enough during the cooler spring and fall weather. As you might guess, there is more to growing blueberries than can be included in a short article. Dr. Art Gaus from the University of Missouri shared this instruction sheet on how to grow blueberries more than 25 years ago. It is still excellent information on blueberry culture. You can access it by going to: <http://www.hfr.ksu.edu/doc3091.ashx>

Blueberries require commitment. Anything less than excellent preparation and care will result in failure. (Ward Upham)

### Blueberries in Containers



Growing blueberries in containers is becoming more popular. Chosen varieties are usually half-high plants that are a cross between highbush and lowbush blueberry species. Plants can be as small as 18 inches tall and wide (Top Hat), but typically are larger. Here are several tips for producing container grown blueberries:

**Acid soil pH:** Blueberries need an acid pH between 4.8 and 5.2. Sphagnum peat moss is very acid and often used in large quantities in soil mixes for acid-loving plants. In fact, blueberries can be grown in peat moss alone if nutrients are provided but that is an involved process. A 50/50 mix of peat moss and potting soil is recommended. This will provide nutrients and weight so the plant is less likely to blow over in wind.

**Container size:** Though containers as small as 2 gallons can be used for half-high blueberries, a larger container will be more stable in the wind and provide a larger moisture reserve during hot, dry weather.

**Watering:** Blueberries do not have root hairs, so they are not efficient in picking up water. Potting soil should be kept moist. This will likely be the most challenging aspect of growing blueberries in containers. A large container will not need to be watered as frequently as a small one.

**Winter care:** Though plants are winter hardy, the roots are not. Move pots into an unheated, attached garage or bury them in the soil or mulch enough to bury the pot in early November. Water them periodically during the winter. Use your finger to determine if the soil is moist one inch deep. If not, then water until some flows out the bottom of the pot.

**Varieties:** Though blueberries will produce some fruit if only a single variety is grown, two varieties will increase the potential fruit crop. Suggested varieties include Top Hat and Northsky. Each should reach about 18 inches high, though Northsky will likely grow wider than Top Hat.

Northblue is another choice that should produce more fruit than either Top Hat or Northsky but should reach 2 to 3 feet high. North Country is intermediate in size at 18 to 24 inches high and should produce a moderate amount of fruit.

**Wind protection:** Wind protection will decrease the amount of water these plants need and reduce the chances of leaf scorch.

## VEGETABLES

### Lettuce



Though lettuce is most often planted directly from seed in late March to early April, it can be started from transplants. Transplants allow lettuce to mature earlier so that it escapes the excessive heat that can lead to a strong flavor and bitterness.

Seed should be started four to five weeks before transplanting. Because transplants are placed at the same time as direct seeding, now would be a good time to begin. Use a seed starting mix and plant shallow as lettuce requires light for germination. A soil media temperature of 60 to 68 degrees will encourage germination. Watch the media temperature carefully, as seed can enter a thermal dormancy if germination temperatures are excessive. Also, a cooler temperature of 55 to 60 degrees should be used once the plants emerge.

Time to maturity varies depending on the type of lettuce, with leaf lettuce being the quickest, followed by bibb, romaine, and buttercrunch lettuce. Head or crisphead lettuce is the slowest and is least likely to mature before becoming bitter.

Spacing also varies with type. Leaf lettuce plants are spaced 4 to 6 inches apart, buttercrunch, bibb, and romaine are set at 6 to 8 inches and head lettuce should be at least 8 inches apart in the row. Lettuce does not have an extensive root system and requires regular watering if rainfall is lacking.

Fertilize before planting according to soil test. Plants should also be sidedressed when about 1/3 grown. Sidedressing is done with fertilizers that have more nitrogen than phosphorus and potassium. Use 1/3 cup of nitrate of soda (16-0-0) or 1/4 cup of a 27-3-3, 29-5-4 or similar fertilizer per 10 feet of row. The latter fertilizers are lawn fertilizers but will work well for sidedressing as long as they do not contain weed killers or weed preventers. (Ward Upham)

**Exposure:** Blueberries do best with a minimum of 6 to 8 hours of sunlight a day. Try a northern or eastern exposure. (Ward Upham)

# TURFGRASS

## Lawn Calendar for Warm-Season Grasses



Following is a lawn calendar for Zoysia-grass and Bermudagrass. Buffalograss, also a warm-season grass, is covered in a separate article.

### March

Spot treat broadleaf weeds if necessary. Treat on a day that is 50 degrees F or warmer. Rain or irrigation within 24 hours of application will reduce effectiveness.

### April

Apply crabgrass preventer between April 1 and April 15, or apply preventer when the eastern redbud is in full bloom. If using a product with proflumicarb (Barricade), apply two weeks earlier. Crabgrass preventers must be watered in before they will start to work.

### May – August 15

Fertilize with 1 lb. of nitrogen per 1,000 square feet per application. Follow the recommendations on the bag. More applications will give a deeper green color, but will increase mowing and lead to thatch buildup with zoysiagrass. Bermudagrass can also have problems with thatch buildup but thatch is less likely with Bermuda than zoysia.

Bermudagrass – Use two to four applications.

Zoysiagrass – Use one to two applications. Too much nitrogen leads to thatch buildup.

One Application: Apply in June.

Two Applications: Apply May and July.

Three Applications: Apply May, June, and early August.

Four Applications: Apply May, June, July, and early August.

### June

If grubs have been a problem in the past, apply a product containing imidacloprid by mid July. Imidacloprid can be applied as early as mid May if there are problems with billbugs or May beetle grubs. These products kill the grubs before they cause damage. They are effective and safe but must be watered in before they become active. June is a good time to core aerate a warm-season lawn. Core aeration will help alleviate compaction, increase the rate of water infiltration, improve soil air exchange and help control thatch.

### Late-July through August

If you see grub damage, apply a grub killer. If Imidacloprid has been applied, this should not be necessary. Grub killers must be watered in immediately.

### Late October

Spray for broadleaf weeds if they are a problem. Treat on a day that is at least 50 degrees F. Rain or irrigation within 24 hours reduces effectiveness. Use the rates listed on the label for all products mentioned. (Ward Upham)

## Lawn Calendar for Buffalograss



### General Comments

Buffalograss has become more popular in recent years due to its reputation as a low-maintenance grass. Buffalograss does require less water and fertilizer than our other turfgrasses but often has problems competing with weeds in eastern Kansas.

Buffalograss is an open growing grass that will not shade the soil as well as most of our other turfgrasses. Weeds are often the result. A regular mowing schedule can reduce broadleaf weed problems as most broadleaves cannot survive consistent mowing. Those that do either have a rosette growing pattern (dandelions, shepherds purse) or are “creepers” (henbit, chickweed, spurge). Annual grasses such as crabgrass or foxtail can also be a problem. A good weed preventer (proflam, pendimethalin or dithiopyr) may be needed prevent problems.

### March

Spot treat broadleaf weeds if necessary. The most important treatment for broadleaf weeds should be in late October to early November well after the buffalograss is dormant. Treatments are much more effective then than in the spring as the weeds are smaller and the weeds are sending energy, as well as the herbicide, to the roots. Treatments in March are to take care of any “escapes” missed in the fall spraying. Spray early enough in March that the buffalograss is still dormant. Look at the base of the plants to make sure there is no green. Treat on a day that is 50 degrees F or warmer. Rain or irrigation within 24 hours of application will reduce effectiveness. Use a combination product such as Trimec, Weed-B-Gon or Weed-Out. Weed Free Zone is also good and will give quicker results under cool conditions.

### April

Apply crabgrass preventer between April 1 and April 15, or apply preventer when the eastern redbud is in full bloom. If using a product with proflam (Barricade), apply two weeks earlier. Crabgrass preventers must be watered in before they will work. Avoid using broadleaf herbicides as the buffalograss is greening up as injury can result. The buffalograss will not be killed but growth will slow making the buffalograss less competitive with weeds.

### June

Fertilize with 1 lb. of nitrogen per 1,000 square feet during June. More applications will give a deeper green color, but will encourage weeds. If it is felt that a second application is needed, apply in July. If grubs have been a problem in the past, apply a product containing imidacloprid by mid July.

Imidacloprid can be applied as early as mid May if there are problems with billbugs or May beetle grubs. These products kill the grubs before they cause damage. They are effective and safe but must be watered in before they become active. Again, I would only treat if grubs have been a problem in the past. Note that the whole area may not need to be treated. The beetles that lay the eggs for the grubs are attracted to lights and moist soil and those areas are most likely to be

infested.

### **Late-July through August**

If you see grub damage, apply a grub killer. If imidacloprid has been applied or if grubs have not been a problem in the past, this should not be necessary. Grub killers must be watered in immediately.

### **Late October to Early November**

Spray for broadleaf weeds if they are a problem. Look carefully as our winter annuals such as chickweed and henbit are small and easily overlooked. Use a product that contains 2,4-D as it increases effectiveness on dandelions. Treat on a day that is at least 50 degrees F. Rain or irrigation within 24 hours reduces effectiveness. Use the rates listed on the label for all products mentioned. (Ward Upham)

## **FLOWERS**

### **Iris Leaf Spot Control Starts Now**



Now is a good time to begin control measures for iris leaf spot by removing old, dead leaves. Iris leaf spot is a fungus disease that attacks the leaves and occasionally the flower stalks and buds of iris. Infection is favored by wet periods during the spring, and emerging leaves eventually show small (1/8- to 1/4-inch diameter) spots. The borders of these spots are reddish, and surrounding tissue first appears water-soaked, and then yellows. Spots enlarge after flowering and may coalesce. The disease tends to be worse in wet weather and may kill individual leaves. Though the disease will not kill the plant directly, repeated attacks can reduce plant vigor so that the iris may die from other stresses. Spores are passed to nearby plants by wind or splashing water.

Because this disease overwinters in old leaves, removal and destruction of dead leaves will help with control. For plants that had little infection the previous year, this may be all that is needed. Plants that were heavily infected last year should be sprayed with chlorothalonil (Bravado Fungicide, Fertilome Broad Spectrum Landscape & Garden Fungicide, Ortho Garden Disease Control, GardenTech Daconil, Bonide Fungonil, Bravo Flowable Fungicide) or myclobutanil (Immunox, Immunox Plus) starting when leaves appear in the spring. Repeat sprays every seven to 10 days for four to six sprays. Iris leaves are waxy, so be sure to include a spreader-sticker in your spray to ensure good coverage. (Ward Upham)

**Contributors:** Ward Upham, Extension Associate

---

To view Upcoming Events: <http://tinyurl.com/fswqe>

The web version includes color images that illustrate subjects discussed. To subscribe to this newsletter electronically, send an e-mail message to [cdipman@ksu.edu](mailto:cdipman@ksu.edu) or [wupham@ksu.edu](mailto:wupham@ksu.edu) listing your e-mail address in the message.

For questions or further information, contact: [wupham@ksu.edu](mailto:wupham@ksu.edu)

*Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.*

*Knowledge  
for Life*

K-State Research and Extension is committed to making its services, activities and programs accessible to all participants. If you have special requirements due to a physical, vision or hearing disability, or a dietary restriction please contact Extension Horticulture at (785) 532-6173.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service K-State Research and Extension is an equal opportunity employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, and United States Department of Agriculture Cooperating, John Floros, Dean.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service K-State Research and Extension is an equal opportunity employer. Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, as amended. Kansas State University, County Extension Councils, and United States Department of Agriculture Cooperating, John Floros, Dean.