

Horticulture 2016 Newsletter

No. 42 October 18, 2016

2021 Throckmorton Plant Science Cntr.
Manhattan, KS 66506 (785) 532-6173

Video of the Week: [Storing Tender Bulbs for the Winter](#)

UPCOMING EVENTS



Kansas Turfgrass Conference in conjunction with KNLA
December 6,7 & 8, 2016



Mark the date to attend the Kansas Turfgrass Conference in conjunction with KNLA on December 6, 7 & 8 in Topeka.

The conference is an excellent way to learn about turf, nursery and landscape management, visit with old friends, network with new ones, and see all the latest equipment and supplies from local and national vendors.

The conference has been approved for Commercial pesticide recertification hours:
1 Core hour 3A - 8 hrs 3B - 9 hrs

International Society of Arboriculture CEUS and GCSAA education points will also be available by attending the conference.

Download a copy of the program, get exhibitor information, or register online
<http://2016ktf.eventbrite.com>

TURFGRASS

Control Broadleaf Weeds in Lawns in Late October - Early November



Late October to early November is the most effective time to control broadleaf weeds in lawns. Dandelions usually produce a flush of new plants in late September, and the winter annual weeds henbit and chickweed should have germinated in October. These young plants are small and easily controlled with herbicides such as 2,4-D or combination products (Trimec, Weed-B-Gon, Weed-Out) that contain 2,4-D, MCPP and Dicamba. Even established dandelions are more easily controlled now than in the

spring because they are actively moving materials from the top portion of the plant to the roots in the fall.

Herbicides will translocate to the roots as well and will kill the plant from the roots up. Choose a day that is 50 degrees or higher. The better the weed is growing, the more weed killer will be moved from the leaves to the roots. Cold temperatures will slow or stop this process.

Weed Free Zone (also sold under the name of Speed Zone) contains the three active ingredients mentioned above, plus carfentrazone. It will give a quicker response than the other products mentioned especially as temperatures approach 50 degrees. (Ward Upham)

Why Late Lawn Seedings Often Fail



We normally recommend that Kentucky bluegrass and tall fescue be seeded in September but no later than October 15. Though plantings later than October 15 can be successful, the odds of success diminish as time passes.

The problem with late plantings is not that the seed will not come up or that young grass plants are sensitive too cold. Most often, the problem is with rooting. Unless the young grass plants have a fairly extensive root system, the freezing and thawing that takes place during winter heaves plants out of the ground, and they dry out and die.

Regardless of when planted, be sure the new lawn is kept watered through the fall. More mature lawns will need less frequent watering but all should go into the winter with moist soil. (Ward Upham)

Tucking Your Lawnmower in for the Winter



If you are done mowing for the year, be sure to service your mower before putting it away. Make sure you drain the gas tank of gasoline-powered engines or use a gasoline stabilizer. Untreated gasoline can become thick and gummy. A few drops of oil squirted inside the spark plug hole (after you remove the spark plug) will help lubricate the cylinder. While you have the spark plug removed, replace it with a new one. If your equipment has a battery, clean the battery terminals, which usually corrode during the season. A wire-bristle brush is a good tool for doing this. The battery can then be removed or connected to a battery monitor that will keep it charged over winter. If you remove the battery, be sure to store it in a protected location for the winter (a cool basement works best). Now is also an excellent time to sharpen mower blades so they'll be ready next spring.

Sharpening rotary mower blades is fairly straightforward. The following steps will guide you through this process:

- * Check the blade for major damage. If you can't fix it, it likely will need to be replaced.

* Remove grass and debris from the blade with a moist cloth. Dry before beginning to sharpen the cutting edge.

* Remove nicks from the cutting edge, using a grinding wheel or hand-file.

* If using a grinding wheel, match the existing edge angle to the wheel. If hand-filing, file at the same angle as the existing edge.

* Grind or file until the edge is 1/32 inch, about the size of a period.

* Particularly with a grinding wheel, avoid overheating the blade as this may warp it.

* Clean the blade with solvent or oil, much like if you were cleaning a gun, for optimum winter storage. Avoid using water because it will promote rust.

Following these tips can help you better prepare your mower for winter storage and also save you some steps this coming spring. (Ward Upham)

VEGETABLES

Hardiness of Cool-Season Vegetables



Cool-season vegetables vary in cold tolerance, with some able to take colder temperatures than others. Semi-hardy crops can take a light frost but are damaged by temperatures in the mid- to upper-20s. Examples include beets, Chinese cabbage, collards, Irish potatoes, Bibb lettuce, mustard, radishes, spinach, Swiss chard, and leaf lettuce. Covering these plants when cold weather threatens can help extend the harvest season.

Plants termed “hardy” can take lower temperatures but are damaged when the temperature drops to the low 20s. These include cabbage, broccoli, cauliflower, Brussels sprouts, carrots, turnips, and kale.

Certain root crops can essentially be stored outside even after the leaves have been damaged or killed by frost. Beets, carrots, potatoes and turnips can be mulched and harvested as needed until the soil starts to freeze in late November to December.

Growing vegetables in Kansas can be a challenge, but we have an extremely long gardening season. We can harvest from early April (asparagus) to early December. Winter is a good time to plan and prepare for next year’s crops. (Ward Upham)

ORNAMENTALS

“Apples” on an Ornamental Shrub

We have had a number of reports of misshapen or fuzzy “apples” developing on a thorny shrub in the landscape. This is most likely flowering quince. Flowering quince is grown for its attractive blooms in the spring. Normally fruit does not develop because frosts prevent fruit formation and plants fruit poorly without



cross-pollination. Quince fruit are quite hard but are sometimes mixed with other fruit to make jellies and preserves. (Ward Upham)

Contributors: Ward Upham, Extension Associate

To view Upcoming Events: <http://hnr.k-state.edu/events/index.html>

The web version includes color images that illustrate subjects discussed. To subscribe to this newsletter electronically, send an e-mail message to cdipman@ksu.edu or wupham@ksu.edu listing your e-mail address in the message.

For questions or further information, contact: wupham@ksu.edu

Brand names appearing in this publication are for product identification purposes only.

No endorsement is intended, nor is criticism implied of similar products not mentioned.

*Knowledge
for Life*

K-State Research and Extension is committed to making its services, activities and programs accessible to all participants. If you have special requirements due to a physical, vision or hearing disability, or a dietary restriction please contact Extension Horticulture at (785) 532-6173.