

Horticulture 2017 Newsletter

No. 7 February 14, 2017

2021 Throckmorton Plant Science Cntr.
Manhattan, KS 66506 (785) 532-6173

Video of the Week:

[How to Make a Grow Light](#)

TURFGRASS

Lawn Calendar for Cool-Season Grasses



The following suggestions are for cool-season grasses such as Kentucky bluegrass or tall fescue. Zoysiagrass, bermudagrass, and buffalograss are warm-season grasses and require a different maintenance regime. A warm-season grass calendar will be covered in a later newsletter.

March

Spot treat broadleaf weeds if necessary. Treat on a day that is 50 degrees or warmer. Rain or irrigation within 24 hours of application will reduce effectiveness.

April

Apply crabgrass preventer when redbud trees are in full bloom, usually in April. The preventer needs to be watered in before it will start to work. One-quarter inch of water will be enough to water in any of the products mentioned in this calendar. Remember that a good, thick lawn is the best weed prevention and may be all that is needed.

May

Fertilize with a slow-release fertilizer if you water your lawn or if you normally receive enough rainfall that your turf doesn't go drought-dormant during the summer. If there are broadleaf weeds, spot treat with a spray or use a fertilizer that includes a weed killer. Rain or irrigation within 24 hours of application will reduce effectiveness of the weed killer, but the fertilizer needs to be watered in. If you are using a product that has both fertilizer and weed killer, wait 24 hours after application before watering in.

June through Mid-July

Apply second round of crabgrass preventer by June 15 – unless you have used Dimension (dithiopyr) or Barricade (prodiamine) for the April application. These two products normally provide season-long control with a single application. Remember to water it in. If grubs have

been a problem in the past, apply a product containing imidacloprid during the first half of July. This works to prevent grub damage. It must be watered in before it becomes active.

Late-July through August

If you see grub damage, apply a grub killer that contains Dylox. Imidacloprid is effective against young grubs but may not be effective on late instar grubs. The grub killer containing Dylox must be watered in within 24 hours or effectiveness drops.

September

Fertilize around Labor Day. This is the most important fertilization of the year. Water in the fertilizer.

November

Fertilize. This fertilizer is taken up by the roots but is not used until the following spring. Water in fertilizer. Spray for broadleaf weeds even if they are small. Broadleaf weeds are much easier to control in the fall than in the spring. Spray on a day that is at least 50 degrees. Rain or irrigation within 24 hours reduces effectiveness. Use label rates for all products! (Ward Upham)

VEGETABLES

Planting Asparagus



Though it is too early to plant asparagus, it is not too early to make plans and prepare soil. This crop is a perennial and will survive for many years if given proper care. It prefers full sun and a well-drained soil and is usually placed on the edge of the garden area so that there is no need to till around the area to plant other crops.

Proper soil prep is especially important for perennial crops. Take a soil test to ensure proper levels of nutrients.

See the accompanying article on how to take a soil test for the correct procedure.

Work the soil as early in the spring as possible but do not work wet soil as clods will form. Then add two inches of organic matter to the surface and the fertilizer and work again so the organic matter and fertilizer are blended into the soil.

Asparagus can be propagated from seed but is more often started from 1-year-old crowns. These crowns are planted deeply; about 8 inches deep either in a hole for each crown or in a trench. Space plants 18 to 24 inches apart. Fill in the trench gradually over the growing season to encourage growth.

March 15 to April 15 is the best planting time. Adapted varieties include Jersey Giant, Jersey King, Jersey Knight, Jersey Supreme and Purple Passion. These are all male hybrids that will produce three times as much as our old Martha or Mary Washington varieties. Males have a number of advantages over females in that they live longer, emerge earlier in the spring, are

more productive and eliminate potential volunteer plants that can reduce the productivity of a planting.

Weed control is very important. Competition with weeds results in slow establishment. A shallow hoeing should be all that is needed. (Ward Upham)

Gardening Calendar

The Kansas Healthy Yards website has a gardening calendar that many gardeners find helpful. It lists all the months of the year and the chores that can be done during that month. Each month is divided into separate sections such as Vegetables and Fruits, Flowers, Lawns, Trees and Shrubs and Houseplants. Not all suggestions will be helpful each year but are important to have included. For example, one of the January tips is on removing snow and ice on limbs. You may brush off the snow, if desired, but ice should be allowed to melt naturally.

To find the gardening calendar, go to <http://kansashealthyyards.org/> and click on “Gardening Calendar” in the gray bar at the top of the page. (Ward Upham)

FLOWERS

Fertilizing Spring-flowering Bulbs



The best time to fertilize spring-flowering bulbs is when foliage emerges in the spring rather than at flowering. Traditionally, gardeners have applied fertilizer during bloom or a bit after, but because bulb roots start to die at flowering, fertilizer applied at bloom is wasted. Roots are active when the foliage first pokes through the ground. Nutrients applied then help the plant produce flowers the following year. If bulbs have been fertilized in the past, there is often plenty of phosphorus and potassium in the

soil. It is best to use a soil test to be certain. If the soil needs phosphorus and potassium, use a complete fertilizer (such as 10-10-10, 9-9-6, etc.) at the rate of 2.5 lbs. per 100 square feet. This would equal 1 rounded teaspoon per square foot. If phosphorus and potassium are not needed, blood meal makes an excellent fertilizer. It should be applied at the rate of 2 lbs. per 100 square feet or 1 teaspoon per square foot. Lawn fertilizers such as a 27-3-3 or 30-3-3 can be used, but cut the rate by a third. Also make sure the lawn fertilizer does not contain a weed preventer or weed killer.

Remember to leave the foliage until it dies naturally. The energy in the foliage is transferred to the bulb as the foliage dies and will help bloom next year. (Ward Upham)

MISCELLANEOUS

Soil Testing



Most gardeners think that soil tests are done only to find out what nutrients are deficient. However, it is just as important to know if you have adequate levels of nutrients so you don't add unneeded fertilizer. The most basic soil test checks pH and the levels of phosphorus and potassium. Most of the lawn and garden soil tests that come out of our soil-testing lab show more than adequate levels of both phosphorus and potassium. If those nutrients are not needed, applying them is a waste of money and can

be a source of pollution. In extreme cases, excess phosphorus can interfere with the uptake of micronutrients. So, if you haven't taken a soil test in several years, take one this spring.

Begin by taking a representative sample from a number of locations in the garden or lawn that goes from the surface to 6 to 8 inches deep. Mix the samples together in a clean container and select about 1 pint of soil. For more detail on taking a soil test, [click here](#).

Take the soil to your local K-State Research and Extension office to have tests done at the K-State soil-testing laboratory for a small fee. A soil test determines fertility problems, not other conditions that may exist such as poor drainage, poor soil structure, soil borne diseases or insects, chemical contaminants or damage, or shade with root competition from other plants (see accompanying article). All of these conditions may reduce plant performance but cannot be evaluated by a soil test. (Ward Upham)

Soil Tests When Soils are Wet



In many parts of the state, this would be a good problem to have. So let's say you just had a rain and need to take a soil test. It would be best to wait until the soil dries but it is possible to take soil tests when soils are wet though there are precautions.

Soil samples should be air-dried before being submitted for testing. Do NOT use artificial means of drying such as an oven or microwave as such treatment may result in inaccurate readings of nutrient levels. Also, be sure to use a clean container to collect the sample. Wet samples are more likely to absorb foreign materials adhering to the container, which may also influence soil test results.

For information on how to take a soil sample, [click here](#).

Take the sample into your local extension office. If you don't know the address for your local, county extension office, see <http://www.ksre.ksu.edu/Map.aspx> (Ward Upham)

Contributors: Ward Upham, Extension Associate

To view Upcoming Events: <http://hnr.k-state.edu/events/index.html>

The web version includes color images that illustrate subjects discussed. To subscribe to this newsletter electronically, send an e-mail message to cdipman@ksu.edu or wupham@ksu.edu listing your e-mail address in the message.

For questions or further information, contact: wupham@ksu.edu

Brand names appearing in this publication are for product identification purposes only.

No endorsement is intended, nor is criticism implied of similar products not mentioned.

*Knowledge
forLife*

K-State Research and Extension is committed to making its services, activities and programs accessible to all participants. If you have special requirements due to a physical, vision or hearing disability, or a dietary restriction please contact Extension Horticulture at (785) 532-6173.