

# Horticulture 2017 Newsletter

No. 4 January 24, 2017

2021 Throckmorton Plant Science Cntr.  
Manhattan, KS 66506 (785) 532-6173

---

**Video of the Week:** [Pothos and Philodendron: Easy Care Plants](#)

## UPCOMING EVENTS



Join us for the [New-Media Marketing Bootcamp](#), February 8-9, 2017 in Overland Park, KS. This event is designed for anyone who plays a role in their brand's social media, from advanced uses to beginners.

Get ideas, inspiration and innovative techniques to implement immediately from sessions including:

- \* Claiming and Building the Online Hub of Your Business: Basic DIY Website Platforms
- \* Learn to Love Your Photos: Clever Phone Photo Tips to Help you Stand Out
- \* Integrating Facebook Ads into Your Marketing Plan
- \* Simple Graphic Tools to Make Your Content Shine
- \* Facebook Deeper Dive: Analyzing Performance of Your Page AND featuring for [Retail Minded](#), Nicole Leinbach Reyhle

More information about [session descriptions](#), [travel information](#), [registration](#) and more can be found on our website menu under New-Media Marketing Bootcamp.

Question? Contact us at [newsmedia@kesu.edu](mailto:newsmedia@kesu.edu)

## VEGETABLES

### Preventing Potatoes from Sprouting in Storage



Home gardeners have had to rely on proper storage conditions (cool and moist) to prevent potatoes from sprouting. But sprouting will eventually occur even if the gardener does everything right. Research by Mary Jo Frazier, Nora Olsen and Gale Kleinkopf from the University of Idaho have found products that should help home gardeners.

These researchers were looking for an organic method to control potato sprouts. They found essential oils from some herbs and spices to be effective sprout inhibitors. Specifically they found that spearmint oil, peppermint oil and clove oil suppressed sprouting by physically damaging rapidly dividing cells in the sprout. Each of these products is so safe that the FDA has approved them for addition to food.

Several application methods were considered though most were only suitable for commercial storage facilities. The only practical method for homeowners was one the researchers labeled a “low-tech” wick method. This was accomplished by placing a small piece of blotter paper saturated with spearmint or peppermint oil in a box with the potatoes. This method was not recommended for the clove oil. Though it was found that peppermint and spearmint oils were equally effective in suppressing sprouts, the peppermint oil was less likely to affect flavor of the potatoes. Reapplication at two- to three-week intervals will be needed for continued sprout suppression. Little to no residue was found on the potatoes from these products due to their high volatility. The first application should be done before sprouting occurs.

Blotting paper is much more difficult to find than it was in the past and so you may want to substitute blank newsprint. However, if blotting paper is desired, try herbarium supply houses. Blotting paper is used to press plant specimens. (Ward Upham)

### Use a Planting Calendar



If you start vegetable plants indoors, it is often helpful to list seeding dates on a calendar so that plants are ready for transplanting at the proper time. To do this, choose your transplant date and count back the number of weeks necessary to grow your own transplants. For example, cabbage, broccoli, and cauliflower are usually transplanted in late March to early April. It takes 8 weeks from seeding to transplant size. Therefore, plants should be seeded in early February.

Information on how many weeks it takes to grow transplants is available in our January 3 newsletter at:

[http://hnr.k-state.edu/extension/info-center/newsletters/2017/January2\\_2017\\_1.pdf](http://hnr.k-state.edu/extension/info-center/newsletters/2017/January2_2017_1.pdf)

Below are examples of some common vegetables grown for transplants and a recommended date for seeding. Dates are Saturdays as this is when many homeowners have the most free time. The dates are not set in stone, and a week earlier or later will not ruin the plants. Also, you may want to seed a week or two earlier if you are in southern Kansas and possibly a week later if you are in northern Kansas. Calendars can be reused year after year by a slight reset of the dates. Also keep notes on how well the transplants did so you can tweak the planting schedule. Your conditions may result in plants that need a bit more or a bit less time.

Crop	<u>Seeding Date</u>	<u>Transplant Date</u>
Cabbage, Broccoli & Cauliflower	February 4	April 1
Lettuce (if you grow		

transplants)	February 4	April 1
Peppers	March 18	May 13
Tomatoes (Ward Upham)	March 25	May 6

## Tomato Trials



Each year we have our Master Gardeners plant and rate a number of tomato varieties. We also give a set to Tom Fowler with the University of Missouri Extension Service. He combines our set with a number of additional varieties. I am still working on data from our Kansas Master Gardeners and will write on those results later. Following are the results from varieties that Tom grew in both 2015 and 2016. All data are from the combined results of both years. We list the average fruit size in ounces, the numbers

of fruit per plant and the number of pounds per plant. I have listed the varieties in alphabetical order.

<b>Variety</b>	<b><u>Avg/Fruit</u></b> <b>(oz)</b>	<b><u>Fruit</u></b> <b><u>Plant</u></b>	<b><u>Lb/</u></b> <b><u>Plant</u></b>
Amana Orange	11.16	4.1	2.98
Amish Paste	2.20	69.6	9.86
Anna Russian	6.42	15.8	6.7
Arkansas Traveler	5.76	19.8	6.32
Beefmaster	8.78	13.8	7.2
Beefsteak	4.51	27.6	6.68
BHN 589	6.08	25	9.48
BHN871	5.43	16.5	5.84
Big Beef	7.34	17.6	5.76
Big Boy	5.25	12.1	4.26
Black Krim	6.26	12	4.64
Brandywine Black	6.86	15.9	6.66
Brandywine Pink	9.31	5.7	3.32
Brandywine Red	8.07	4.7	2.22
Brandywine Yellow	3.74	0.6	0.28
Celebrity	5.45	25.5	8.7
Chef's Choice	5.88	25.9	9.52
Cherokee Purple	7.95	13.4	6.76
Delicious	3.02	18.6	4.2
Early Girl	3.17	31.7	6.1
Florida 91	6.33	19	7.52
German Johnson	7.72	4.1	1.95
Giant Belgium	9.27	7.3	4.6
Golden Jubilee	5.20	8	2.35
Jet Star	5.78	31.4	11.1
Mortgage Lifter	7.36	7.5	3.52
Mt. Majesty	6.19	25.8	10.14
Oxacan Jewel	4.58	12.4	3.6
Oxheart	5.35	14.9	4.98
Ponderosa Pink	7.84	6	2.95
Roma	1.78	51.7	5.78
Rutgers	2.91	22.3	4.12
Summer Pick	5.88	22.4	8.6
Ultrasonic	5.00	32.7	10.44
Vintage Wine	6.50	3.5	1.44

Both 2015 and 2016 were challenging years for northwest Missouri with 2016 being the most difficult. It will be interesting to compare what Tom found with the results from our Master Gardeners. (Ward Upham)

## FRUIT

### Cloning Apple Trees



We occasionally receive calls from gardeners who want to know how to treat an apple seed so it will germinate. Usually, the gardener is trying to reproduce an old apple tree that was special for some reason (good quality fruit, planted by grandparents, etc.) Unfortunately, apples grown from seed will not be like the parent. About 1 in every 80,000 apple trees grown from seed will be as good as the apples we are used to eating. Apple trees grown from seed usually have small and inferior quality fruit.

If you want a tree exactly like the parent, you must propagate that tree vegetatively. In the case of apples, this usually means grafting. Apple trees are actually quite easy to graft, even for novices. Don't be afraid to try even if you haven't grafted before. The step that needs to be done at this time of year is the choosing and cutting of scion wood or small branches that will be grafted on top of a rootstock.

See the accompanying article in this newsletter on how this is done. However, if you don't have an existing tree to graft onto, you will need to plant a rootstock this year for grafting onto next. Fruit trees are normally grafted (or budded) onto specially selected rootstocks. These rootstocks usually reduce tree size. For example, a tree that normally would reach 25 feet tall will only reach 10 feet if it is grown on a certain rootstock. Dwarfing rootstocks also allow apples to bear fruit a year or more earlier.

A tree on its own roots normally takes 5 to 7 years before it will bear. Semi-dwarf trees bear in 4 to 5 years, and dwarf trees bear in 3 to 4 years. Unfortunately, not all dwarfing rootstocks are well adapted to Kansas. Semi-dwarf trees usually are a better choice for us. Note that rootstock reduces tree size, not fruit size. Therefore, a Golden Delicious tree that only reaches 8 feet tall due to a dwarfing rootstock, will bear the same size fruit as a Golden Delicious tree that is 25 feet tall.

Most nurseries only sell trees that are already grafted. A company that does sell rootstocks is Raintree Nursery, Morton, WA, (360) 496-6400, <http://www.raintreenursery.com/Rootstocks/> Another is Cummins Nursery, (865) 233-3539, <http://www.cumminsnursery.com/rootstocks.htm> though there is a minimum shipping and handling fee of \$20.

It is also possible to buy a tree from a local nursery and graft your clone into it. This will give you one tree that produces two different apples. One disadvantage of this method is that it is possible to prune off the special clone by mistake in later years.

This information does not include the details of grafting or budding or subsequent care. The Missouri Extension Service has an excellent publication on grafting at <http://extension.missouri.edu/explorepdf/agguides/hort/g06971.pdf> as well as a second publication on budding at <http://extension.missouri.edu/explorepdf/agguides/hort/g06972.pdf> . Be sure to practice your cuts on wood you prune off in March. After you do about 100 cuts, you will start to get the hang of it. (Ward Upham)

## Multiple Grafts on Apple Trees



Nurseries often sell apple trees that bear more than one variety of fruit. The secret is grafting. All apple trees are grafted, which is done by grafting the apple-producing variety (the scion) on a variety chosen for its dwarfing effects (the rootstock). A tree with more than one variety simply has more than one fruiting variety grafted onto a single rootstock. Grafting allows growers to have a single tree that could produce Jonathan, Red Delicious and Golden Delicious apples. These trees can be a unique attraction and a good conversation point in a fruit garden. If space is limited, a multiple grafted tree may allow growers to have a greater variety of fruit than with individual varieties on separate trees. However, there are some possible drawbacks. Whoever prunes the trees may not recognize the individual grafts and may unknowingly prune off one of the varieties. Also, varieties may vary in vigor, and stronger varieties can crowd weaker ones. There also may be a difference in susceptibility to disease among varieties and among different kinds of fruit. Some may have resistance to a disease and not require protection, but others are susceptible and do require protection. If the susceptible fruits are protected, the more resistant ones will be sprayed unnecessarily. (Ward Upham)

## MISCELLANEOUS

### Growing Your Own Firewood



In order to avoid energy costs, some homeowners are turning to wood for heat. Plant species is an important consideration as not all trees have the same density and therefore, heat value. The greater the dry weight, the better.

The highest value for trees commonly found in Kansas is osage orange (hedgeball tree) at 4,800 pounds per cord. Osage orange has a gnarly growth habit and a nasty set of thorns. This species also sparks which isn't a problem in a wood-fired boiler but certainly would be in an open fireplace.

Black locust is next with 4,200 pounds per cord. Black locust is a fast grower and also has excellent burning qualities and makes a nice bed of coals. However, it is hard to split, suckers, and has some relatively small thorns, especially on young trees.

Buroak and red oak come in at 3,800 and 3,500 pounds per cord respectively but are not fast growers. Mulberry, however, has the same weight as red oak but grows more quickly. Silver maple has less heat value (3,000 pounds per cord) but is a very fast growing tree.

Black locust would be my first choice for this purpose though you may wish to plant rows of several species. However, each situation is different and another species may work better for you. So how do you set out your plantation? Dr. Wayne Geyer, our late forestry professor, did many woody biomass studies over a period of 35 years. Following are some recommendations that have come out of his studies.

- Plant locust a few rows in from a field edge to reduce suckering in the field.
- Plant on a close spacing, 4 to 6 feet apart. This maximizes yield and reduces side branching.
- Control weeds the first two years.
- Harvest every 5 years, most trees will resprout and can be reharvested.
- Plant about 1 acre per year for 5 years if you wish to supply the majority of the firewood needed to heat your home.

Trees mentioned above and available from the Kansas Forest Service include osage orange, bur oak, red oak and silver maple. (Ward Upham)

---

To view Upcoming Events: <http://hnr.k-state.edu/events/index.html>

The web version includes color images that illustrate subjects discussed. To subscribe to this newsletter electronically, send an e-mail message to [cdipman@ksu.edu](mailto:cdipman@ksu.edu) or [wupham@ksu.edu](mailto:wupham@ksu.edu) listing your e-mail address in the message.

For questions or further information, contact: [wupham@ksu.edu](mailto:wupham@ksu.edu)

*Brand names appearing in this publication are for product identification purposes only.*

*No endorsement is intended, nor is criticism implied of similar products not mentioned.*

*Knowledge  
for Life*

K-State Research and Extension is committed to making its services, activities and programs accessible to all participants. If you have special requirements due to a physical, vision or hearing disability, or a dietary restriction please contact Extension Horticulture at (785) 532-6173.