

Problem: Blister Beetles



Hosts: Tomatoes, beans, peas, potatoes, and other vegetables.

Description: Though there are several members of this family that can cause damage, the Striped and Gray Blister Beetles are the most common. The adult stage causes the damage; not the larvae. The beetles tend to move in swarms and can cause a great deal of defoliation but may not stay in one area for very long.

Blister beetles pass the winter in a pseudopupa stage and then go through a final molt in the spring. After a short period of activity, the larva enters the true pupal stage with the adults emerging midsummer.

Blister beetles have long, slender bodies with a relatively large head. These insects release a caustic substance when crushed that can raise blisters on the skin.

Recommendations: Often beetles will move on in a day or two. If beetles are handpicked, be sure to wear gloves. A number of stomach insecticides such as cyfluthrin (BioAdvanced Vegetable & Garden Insect Spray), permethrin (Bonide Eight Vegetable, Fruit and Flower Concentrate and Hi-Yield Garden and Farm Insect Control) and gamma-cyhalothrin (Spectracide Triazicide) are also effective for control.

References:

1. Insect Pests, A Golden Guide, Golden Press, pg 74.
2. Garden Insects of North America, Princeton University Press, pg 176

Last Update: 1/4/2022

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

“Knowledge for Life”

Kansas State University Agricultural Experiment Station and Cooperative Extension Service