

Problem: Rhubarb Seedstalks



Host Plants: Rhubarb

Description: Rhubarb is a perennial vegetable that produces leaves early in the season that are harvested for the petiole (leaf stalk). Later in the season- especially in seasons with a longer, cool spring season, the plant will try to produce a large, hollow-stemmed seedstalk from the center of the plant. If the rhubarb is allowed to flower and produce seed, the vigor of the plant is reduced.

Recommendations: Seedstalks should be removed as soon as they appear. It may seem like a losing battle since seedstalks may rapidly re-appear. Keep cutting them out as they form and the seedstalk development will quit in a few weeks. Rhubarb does not come 'true to type' from seed so there is no advantage in allowing the plant to produce seed and collect the seed for re-planting. Most improved strains of rhubarb are selected for their bright red-colored stalks, vigor, and tendency to produce few, if any, seedstalks.

References:

1. [Rhubarb](#), Kansas State University, KSU Horticulture Report EP99
2. [Grow Your Own Rhubarb](#), Oregon State University, EC 797

Last Updated: 11/13/2023

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

“Knowledge for Life”

Kansas State University Agricultural Experiment Station and Cooperative Extension Service