

Problem: Blossom-end Rot



Plants Affected: Tomato, Pepper, Squash, Eggplant and Watermelon.

Description:

This condition is easily recognized by the flat, leathery area on the blossom end of the fruit - opposite the stem end. This condition is often first seen when the fruit is about half-grown and continues as it matures. This is not a disease but a non-parasitic disorder that is most common on tomatoes.

Blossom-end rot is the result of a calcium deficiency in the blossom-end of the fruit. Calcium is required in relatively large amounts when fruit is growing rapidly. If demand exceeds supply, deprived tissues break down leaving the characteristic leathery bottom. This may be the result of low calcium levels in the soil (rare in Kansas) but can also be caused by a sharp change from cool to hot weather, extreme soil moisture fluctuations, drought stress, root damage due to deep cultivation or waterlogged conditions, or excessive growth due to too much nitrogen fertilization. Blossom-end rot due to less than adequate uptake of calcium because of a limited root system will correct itself as the plant compensates by adding more roots.

Recommendations:

1. Maintain soil pH around 6.5 as shown by soil test. Do **not** assume that blossom-end rot is sure evidence that the soil lacks calcium. Other conditions that interfere with calcium uptake can cause calcium to be demanded in higher quantities than the plant can physically supply.
2. Use a nitrate nitrogen rather than one that releases nitrogen in the ammonium form. Excess ammonium ions reduce calcium uptake. Also make sure that the plants are not overfertilized especially when the fruit are small. Usually 1½ pounds of a 10-20-10 fertilizer applied before planting is sufficient for tomatoes though sandy soils may need monthly sidedressings.

3. Maintain even soil moisture. Mulches conserve moisture and moderate soil moisture extremes. Even with mulches, weekly irrigation will be necessary if there is no rain. Fluctuations in soil moisture often leads to an increase in blossom-end rot.
4. Make sure tomatoes are planted in an area with good drainage. Excess water causes roots to smother leading to reduced calcium uptake.
5. Do not hoe too deeply. Deep hoeing can damage roots and thereby reduce calcium uptake.

Blossom-end rot can occur even to well cared for plants if weather conditions are favorable. Fortunately, the condition normally corrects itself in a couple of weeks. Gardeners may wish to remove damaged tomatoes to allow more energy to flow to the remaining unaffected fruit.

References:

1. [Blossom-End Rot of Tomato, Pepper, and Eggplant](#), Ohio State University Extension FactSheet (HYG-3117-96)
2. [Blossom-end Rot of Tomatoes](#), Oregon State University Extension Publication (FS 139)
3. [Growing Tomatoes in the Home Garden](#), North Carolina State University Leaflet (Leaflet No: 28-D)

Last Update: 11/20/2015

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

“Knowledge for Life”

Kansas State University Agricultural Experiment Station and Cooperative Extension Service