

Problem: Brownheaded Ash Sawfly - *Tomostethus multicinctus*



Hosts: Green ash is most commonly attacked but other ash are susceptible.

Description: These sawfly larvae are a light green color with a broad, whitish stripe on the top side. In the middle of the whitish stripe there appears to be a darker green stripe that is actually the digestive tract of the insect. Though sawfly larvae resemble caterpillars, they have at least six pairs of "stublike" prolegs behind the three pairs of true legs on their abdomen. (Caterpillars never have more than five pairs of prolegs.)

Recommendations: There are usually no detrimental effects to the health of the tree if nature is allowed to run its course. Even if all the leaves are eaten, damage is early enough in the growing season for trees to put out a complete new set of leaves and have enough time to make all the food reserves needed to survive the coming winter. However, if control is desired, a number of insecticides can be used for including cyfluthrin (Bayer Vegetable & Garden Insect Spray), malathion, esfenvalerate (Monterey Bug Buster II) and carbatyl (Sevin). An effective organic product is spinosad (Captain Jack's Deadbug Brew; Fertlome Borer, Bagworm, Leafminer and Tent Caterpillar Spray, Monterey Garden Insect Spray). Horticultural oils and insecticidal soaps are also effective because of the soft skin of sawfly larvae.

References:

1. Garden Insects of North America, Whitney Cranshaw, pg 78
2. [Ash Sawfly](#), Plantalk Colorado, #1455

Last Update: 1/11/2018

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

“Knowledge for Life”

Kansas State University Agricultural Experiment Station and Cooperative Extension Service