

Problem: Bacterial Spot of Peach - *Xanthomonas compestris* pv *pruni*



Host Plants: Peach and nectarine and, to a lesser extent, apricot and plum

Description: This bacterial disease is a common disease in Kansas but rarely causes extensive losses in fruit quality or production. The disease affects leaves, fruit, and occasionally twigs. Leaf infections first appear as small, pale green to yellow angular spots. These spots normally are more prevalent near the leaf tip and are surrounded by a yellow halo. Eventually, the lesions turn dark brown, separate from healthy tissue, and drop out, giving the leaf a shot-hole appearance. On susceptible cultivars, 1 or 2 leaf lesions can cause a complete yellowing of the leaf and result in premature defoliation.

One of the most serious aspects of the disease is the development of spots on the fruit. The spots are mostly superficial, but numerous infections can cause pitting and cracking of the fruit surface. The fruit cracks may show gumming.

The bacterium overwinters in small, blister-like cankers on the twigs. In early spring, the cankers rupture and release millions of bacteria. These bacteria are dispersed by splashing water and infect developing leaves and fruit. Infection is favored by frequent rainfall accompanied by high winds and moderate temperatures. Extended periods of hot, dry weather inhibit disease development.

Recommendations: Peach cultivars vary in their susceptibility to bacterial spot, although no variety is immune. Those cultivars showing tolerance to the disease include Belle of Georgia, Biscoe, Candor, Comanche, Dixired, Earliglo, Early-Free Red, Emery, Encore, Garnet Beauty, Harbelle, Harbinger, Harbrite, Harken, Late Sunhaven, Loring, Madison, Norman, Ranger, Redhaven, Redkist, Redskin, Sentinel, and Sunhaven.

Cultivars most susceptible to bacterial spot are Autumnnglo, Autumn Lady, Blake, Elberta, Halehaven, July Elberta, Jersey Queen, Jerseyland, Kalhaven, Suncling, Suncrest, Sunhigh, Ran Cocas, Redcrest, Rio-Oso-Gem, and Sweet Sue. Most apricot and nectarine cultivars are susceptible to bacterial spot.

Excessive fertilization can predispose peaches to severe bacterial spot infections. On the other hand, proper fertilization is necessary to maintain tree vigor and to prevent the development of Cytospora canker. Monitoring the fertility needs of trees is best accomplished through soil and leaf analysis.

References:

1. [Bacterial Spot of Peach](#), Plant Disease Diagnostic Clinic Factsheet, Cornell University

Last Update: 10/11/2023

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

“Knowledge for Life”

Kansas State University Agricultural Experiment Station and Cooperative Extension
Service