

**Problem:** Tan or White Drupelets**Plants Affected:** Blackberry and raspberry

**Description:** Blackberry and raspberry fruit will often develop white or tan drupelets on the berry. Though we are not completely sure of the cause, two commonly given reasons are stinkbug damage and sunscald. Damage has been attributed to stinkbugs if the pattern of off color (not white) drupelets is random. Stinkbug damage is caused by the insect feeding on the blackberry receptacle and injuring drupelets on either side. Sunscald damage will be on the side of the fruit exposed to the sun and has several drupelets in a small area being affected.

**Recommendations:** Neither condition affects the eating quality of the fruit unless the stink bug releases the “stink” with which it is associated rendering the fruit inedible. By the time damage is seen, it is too late for control.

**References:**

1. [Other Notable Blackberry Insects – Harvest Pests](#), NC Small Fruit, Specialty Crop, and Tobacco IPM, June 29, 2010
2. [The Problem of White Drupelets in Blackberries](#), Mississippi Fruit and Nut Blog, June 18, 2015

**Last Update:** 1/11/2018

---

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

*“Knowledge for Life”*

Kansas State University Agricultural Experiment Station and Cooperative Extension Service