

Problem: Woolly Elm Aphid (*Eriosoma americana*)



Hosts: Elm

Description: Leaves are twisted and malformed with leaf clusters forming rosettes. Unfurling these leaves reveals aphids that are covered with white waxy threads.

Life History: The Woolly Elm Aphid overwinters as an egg placed in a bark crevice. These eggs are produced by females the previous fall. It is interesting to note that these females produce a single egg and die soon after the egg is laid because they lack functional mouth parts. Eggs hatch in the spring producing a wingless female that feeds by sucking sap from the bottom of a young elm leaf.

At maturity, these females produce nearly 200 young, which are, again, all females. Leaves form rosettes as populations rise. A winged generation (all females) appears by the end of June and actively seeks out the Amelanchier shrub which is commonly known as the shadbush or serviceberry. A new generation of females migrates to the underground parts of the shrub. Several generations are produced underground. In the fall, a final winged generation of females migrates back to the elm and starts the cycle over again.

Recommendations: Damage is usually minimal and control is difficult due to protection provided to the aphids by the rosette of leaves. Control measures are rarely warranted.

References:

1. [Aphids in the Home Yard and Garden](#), University of Minnesota Extension
2. *Insects That Feed on Trees and Shrubs*, Cornell University Press, pg 262.

Last Update: 1/11/2022

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

“Knowledge for Life”

Kansas State University Agricultural Experiment Station and Cooperative Extension Service