

# Horticulture 2021 Newsletter

No. 8 February 23, 2021

---

1712 Claflin, 2021 Throckmorton Plant Science Cntr.  
Manhattan, KS 66506 (785) 532-6173

**Video of the Week:** [Easy to Grow Peas](#)

## REMINDERS

1. Prune fruit trees
2. Check fruit trees for scale and apply dormant oil if present..

## VEGETABLES

### Cure the Itch by Planting Peas



If you are tired of winter and hunger for spring, try planting peas as soon as the soil dries and the soil temperature reaches 40 degrees. There are several types of peas we can plant in Kansas. Probably the most common is the shelling pea and the old standard in this group is Little Marvel. Though Little Marvel is still on our recommended list, we have a number of others that do well including Green Arrow, Knight, Maestro, Burpeeana and Mr. Big. All of these are early maturing types that allow us to harvest a crop before the hot weather arrives and stops production.

Snow peas are those commonly used in stir-fry that have a crisp edible pod. Recommended varieties include Dwarf Grey Sugar and Mammoth Melting Sugar.

Sugar snap peas resemble shelling peas but have a thick, fleshy pod and can be eaten fresh, steamed or cooked. Like snow peas, they are not shelled but eaten pod and all. We recommend Sugar Bon, Sugar Ann, Super Sugar Snap and Sugar Sprint.

Peas should be planted shallow, about one-half inch deep, to encourage rapid germination and emergence. Seed in the row should be spaced 2 inches apart. Many people often plant two rows 6 to 8 inches apart so the floppy plants can support one another. For some older varieties, this may not be enough. They may need trellising to support the growing vines. Fencing may be needed to keep rabbits away. (Ward Upham)

### Lettuce



Though lettuce is most often planted directly from seed in late March to early April, it can be started from transplants. Transplants allow lettuce to mature earlier so that it escapes the excessive heat that can lead to a strong flavor and bitterness.

Seed should be started four to five weeks before transplanting. Because transplants are planted at the same time as direct seeding, now would be a good time to begin. Use a seed starting mix and plant shallow as lettuce requires light for germination. A soil media temperature of 60 to 68 degrees will encourage germination. Watch the media

temperature carefully, as seed can enter a thermal dormancy if germination temperatures are excessive. Also, a cooler temperature of 55 to 60 degrees should be used once the plants emerge.

Time to maturity varies depending on the type of lettuce, with leaf lettuce being the quickest, followed by bibb, romaine, and buttercrunch lettuce. Head or crisphead lettuce is the slowest and is least likely to mature before becoming bitter.

Spacing also varies with type. Leaf lettuce plants are spaced 4 to 6 inches apart, buttercrunch, bibb, and romaine are set at 6 to 8 inches and head lettuce should be at least 8 inches apart in the row. Lettuce does not have an extensive root system and requires regular watering if rainfall is lacking.

Fertilize before planting according to soil test. Plants should also be sidedressed when about 1/3 grown. Sidedressing is done with fertilizers that have more nitrogen than phosphorus and potassium. Use 1/3 cup of nitrate of soda (16-0-0) or 1/4 cup of a 27-3-3, 29-5-4 or similar fertilizer per 10 feet of row. The latter fertilizers are lawn fertilizers but will work well for sidedressing as long as they do not contain weed killers or weed preventers. (Ward Upham)

### Soil Temperature and Vegetables



One of the most neglected tools for vegetable gardeners is a soil thermometer. Soil temperature is a much better measure of when to plant than air temperature or the calendar. Planting when soil is too cool can cause some seeds to rot and transplants to sit there.

A number of vegetables can germinate and grow at cool temperatures. For example, peas will germinate and grow well at a soil temperature of 40 F. Though lettuce, parsnips, and spinach can sprout at a soil temperature of 35 F, they prefer at least 45 F for best germination and growth. Radishes also do well at a soil temperature of 45 F. Even if the seeds of these cool-season crops are planted below the recommended soil temperature, the seed will rarely rot.

Warm-season crops such as tomatoes, sweet corn and beans are different. They prefer at least 55 F for germination (or transplanting), but others such as peppers, cucumbers, melons and sweet potatoes need it even warmer, about 60 F. If planted when soils are too cool, they likely will rot before germinating.

Taking soil temperature accurately is a bit of a science. First, use a thermometer with a metal probe. These are sold in many garden, auto parts and hardware stores. Those in auto parts stores are used to measure the temperature inside air conditioning ducts and are often less expensive than those used for gardening. Take the temperature 2.5 inches deep at about 10 to 11 a.m. Temperature variations throughout the day and night affect soil temperature, with lowest readings after dawn and warmest around mid-afternoon. The late-morning reading gives a good average temperature. If taking the soil temperature at this time is not practical, take a reading before you leave for work and a second when you return home and use the average. Also be sure to get a consistent reading for four to five days in a row before planting, and make sure a cold snap is not predicted.

An excellent guide sheet on this subject is published by the Alabama Cooperative Extension System and is titled "Soil Temperature Conditions for Vegetable Seed Germination." It can be found at <https://tinyurl.com/1jw297zt> (Ward Upham)

# HOUSEPLANTS

## Leaching Houseplants



Everyone knows that someone stranded in the ocean should not drink the water. The salt content of that water will make a bad situation worse. What many people don't realize is that this same principle can harm plants.

Fertilizers are salts or are converted to salts before plant take up. They must be salts in order for the plant roots to absorb them. However, salt levels can build up over time and eventually may harm plant roots leading to scorched leaves and unhealthy plants. Though this can happen under field conditions,

especially in low rainfall areas, it is particularly critical with houseplants.

Houseplants have a certain soil volume that doesn't change until a plant is repotted. Salt build-up can be a crucial concern especially if the houseplants are fertilized so heavily that the plants can't use all the nutrients and fertilizer salts build up. This is especially common in winter when houseplants do not use much fertilizer due to low levels of light.

Leaching an overabundance of salts can be an important practice to ensure the health of our houseplants. Leaching is not a complicated or difficult process. It consists of adding enough water to wash out excess salts.

How much water is enough? Add the amount of water that would equal twice the volume of the pot. This, of course, would need to be done outside or in a bathtub or sink. Water must be added slowly so that it doesn't overflow the rim of the pot. If salt has formed a crust on the surface of the soil, remove it but don't take more than 1/4 inch of the underlying media. This may also be a good time to repot the plant. (Ward Upham)

## An Easy Way to Propagate House Plants



You don't need a lot of equipment to propagate a houseplant. Gardeners can get by with a coffee cup, potting soil, 3 drinking straws, a plastic bag and a rubber band. Start by making a slit or hole in the bottom of the coffee cup so that it drains excess water. Then fill the cup with moist potting soil. Do not use garden soil as it does not drain well. Too much water (and too little oxygen) will harm cuttings.

### *Prepare the Cutting*

- Remove about a 4-inch or smaller piece from the tip of the plant. The cut should be made just below a node. A node is where a leaf attaches to the stem.
- Remove the leaf or leaves from the bottom node. This is where roots will form.
- If there are just a few leaves on the tip, fine. However, if there is a cluster of leaves, remove most of them below the tip. This will cut down on water loss as the plant makes new roots.

### *Plant the Cutting*

- Push the bottom end of the cutting into the soil. The remaining leaves should not contact the soil. A rooting hormone may be used if desired but usually is unnecessary with houseplants.

### *Make a Greenhouse*

- Place 3 straws equidistant from each other near the outside edge the cup full of potting soil. They will

support the plastic bag so that it does not contact the leaves and cause them to rot.

- Place the plastic bag over the cup like a tent and use the rubber band to secure the open end of the bag to the sides of the cup.

#### *Grow the Cutting*

- Place the cutting in bright, indirect light. Do not place in full sunlight as the cutting may overheat.

- Keep the cutting warm. A temperature of 72 degrees is ideal. Roots should form in about 10 days.

Check by removing the plastic bag and pulling gently on the cutting. If it doesn't pull out easily, roots have started to form and the plastic bag can be left off. (Ward Upham)

## FLOWERS

### **Iris Leaf Spot Control Starts Now**




Now is a good time to begin control measures for iris leaf spot by removing old, dead leaves. Iris leaf spot is a fungus disease that attacks the leaves and occasionally the flower stalks and buds of iris. Infection is favored by wet periods during the spring, and emerging leaves eventually show small (1/8- to 1/4-inch diameter) spots. The borders of these spots are reddish, and surrounding tissue first appears water-soaked, and then yellows. Spots enlarge after flowering and may coalesce. The disease tends to be worse in wet weather and may kill individual leaves. Though the disease will not kill the plant directly, repeated attacks can reduce plant vigor so that the iris may die from other stresses. Spores are passed to nearby plants by wind or splashing water.

Because this disease overwinters in old leaves, removal and destruction of dead leaves will help with control. For plants that had little infection the previous year, this may be all that is needed. Plants that were heavily infected last year should be sprayed with chlorothalonil (Bravo Fungicide, Fertilome Broad Spectrum Landscape & Garden Fungicide, Ortho Garden Disease Control, GardenTech Daconil, Bonide Fungonil, Bravo Flowable Fungicide) or myclobutanil (Immunox, Fungi-Max, Fertilome F-Stop Lawn & Garden Fungicide) starting when leaves appear in the spring. Repeat sprays every seven to 10 days for four to six sprays. Iris leaves are waxy, so be sure to include a spreader-sticker in your spray to ensure good coverage. (Ward Upham)


## MISCELLANEOUS

### **Pruning Deciduous Shrubs**

Before Pruning



After Pruning



Gardeners are eager to get out and do something in the landscape this time of year. One chore that can be taken care of now is pruning certain shrubs. Often, gardeners approach pruning with trepidation, but it is not as difficult as it may seem. Remember, not all shrubs need to be pruned (i.e. witch hazel), and certain

shrubs, which will be identified later, should not be pruned this time of year. Shrubs are pruned to maintain or reduce size, rejuvenate growth, or to remove diseased, dead or damaged branches. Deciduous shrubs are those that lose their leaves each winter. Evergreen shrubs maintain foliage all year and include yews and junipers.

Deciduous shrubs are placed into three groups:

- Those that flower in the spring on wood produced last year;
- Those that flower later in the year on current season's growth; and
- Those that may produce flowers, but those flowers are of little ornamental value.

Shrubs that flower in the spring should not be pruned until immediately after flowering. Though pruning earlier will not harm the health of the plant, the flowering display will be reduced. Examples of these types of plants include forsythia, lilac and mock orange. Shrubs that bloom on current season's growth or that do not produce ornamental flowers are best pruned in late winter to early spring. Examples include Rose-of-Sharon, pyracantha, Bumald spirea and Japanese spirea.

Pruning during the spring allows wounds to heal quickly without threat from insects or disease. There is no need to treat pruning cuts with paints or sealers. In fact, some of these products may slow healing. There are three basic methods used in pruning shrubs: thinning, heading back and rejuvenating. Thinning is used to thin out branches from a shrub that is too dense. It is accomplished by removing most of the inward growing twigs by cutting them back to a larger branch. On multi-stemmed shrubs, the oldest canes may be completely removed.

Heading back is done by removing the end of a branch by cutting it back to a bud and is used for either reducing height or keeping a shrub compact. Branches are not cut back to a uniform height because this results in a "witches-broom" effect.

Rejuvenation is the most severe type of pruning and may be used on multi-stem shrubs that have become too large, with too many old branches to justify saving the younger canes. All stems are cut back to 3- to 5-inch stubs. This is not recommended for all shrubs but does work well for spirea, forsythia, pyracantha, ninebark, Russian almond, little leaf mock orange, shrub roses and flowering quince. (Ward Upham)

### **Did the Extreme Cold Hurt This Year's Fruit Crop?**



Normally we are concerned with cold damage to fruit buds as they break dormancy in the spring. However, this year, people are concerned that the extreme cold temperatures last week may have damaged their fruit plants. Though most fruit plants should have come through the cold temperatures without significant damage, fruit buds or fruit canes are more sensitive. For example, thornless blackberry canes may have been killed to the ground. However, the thornless blackberry plant is likely fine and will put up new growth this summer. Raspberries and thorny blackberry fruiting canes may have survived but it will be a wait and see proposition.

What about tree fruit? The trees are likely fine but fruit buds may have been affected. Usually we start losing the fruiting buds of peaches and nectarines when the temperature drops to 10 degrees F below zero. Cherry buds are more hardy and will often withstand temperatures down to -15F. Apple and pear are even more hardy and often escape damage until the temperature reaches -25 to -30. However, there is significant variability among different varieties or even different localities and you may receive damage at temperatures less severe than those mentioned. Also, you may not find damage even though temperatures dropped below those listed. (Ward Upham)

**Contributors:** Ward Upham, Extension Associate

---

Division of Horticulture  
1712 Claflin, 2021 Throckmorton  
Manhattan, KS 66506  
(785) 532-6173

For questions or further information, contact: [wupham@ksu.edu](mailto:wupham@ksu.edu) OR [cdipman@ksu.edu](mailto:cdipman@ksu.edu)

This newsletter is also available on the World Wide Web at:

<http://hnr.k-state.edu/extension/info-center/newsletters/index.html>

The web version includes color images that illustrate subjects discussed. To subscribe to this newsletter electronically, send an e-mail message to [cdipman@ksu.edu](mailto:cdipman@ksu.edu) or [wupham@ksu.edu](mailto:wupham@ksu.edu) listing your e-mail address in the message.

Brand names appearing in this newsletter are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

K-State Research and Extension is committed to making its services, activities and programs accessible to all participants. If you have special requirements due to a physical, vision or hearing disability, or a dietary restriction please contact Extension Horticulture at (785) 532-6173.